

## Effect of respiratory muscle exercise on the improvement of respiratory function in asthmatic patients; Systematic review

Mohammed Sagheer Albarqi <sup>1,\*</sup>, Hussein Saleh Alyami <sup>2</sup>, Rakan Abdullah Alshareef <sup>1</sup>, Saad alsaad <sup>3</sup>, Fahad Bader AlGhounaim <sup>4</sup>, KAMAL KAMEL ALSOYANI <sup>1</sup>, Mohammed Sultan Alshehri <sup>1</sup>, Mortada Hassan Aljassas <sup>1</sup>, Mania Salem Al-baqawi <sup>3</sup>, Ahmad Abdulrazak Al Dulaijan <sup>5</sup> and Abdullah Mohammed Al-Mutairi <sup>6</sup>

<sup>1</sup> Department of Respiratory Therapy, Imam Abdulrahman Bin Faisal Hospital, NGHA, Dammam, Saudi Arabia.

<sup>2</sup> Imam Abdulrahman Bin Faisal Hospital, NGHA, Dammam, Saudi Arabia.

<sup>3</sup> Department of Respiratory Therapy, Imam Abdulrahman Alfaaisal Hospital, NGHA, Dammam, Saudi Arabia.

<sup>4</sup> Department of Home Health Care, King Abdulaziz Hospital, NGHA, Al Ahsa, Saudi Arabia.

<sup>5</sup> Emergency Nurse, Emergency Department, Imam Abdulrahman Bin Faisal Hospital, NGHA, Dammam, Saudi Arabia.

<sup>6</sup> Anesthesia technician, Anesthesia department, Imam Abdulrahman Alfaaisal, NGHA, Dammam, Saudi Arabia.

World Journal of Advanced Research and Reviews, 2019, 03(01), 036-042

Publication history: Received on 12 July 2019; revised on 26 August 2019; accepted on 29 August 2019

Article DOI: <https://doi.org/10.30574/wjarr.2019.3.1.0042>

### Abstract

**Background:** Comprehensive programs that combine medication-assisted asthma treatment with education, breathing techniques, and fitness training have been highlighted. Examining the effects of respiratory muscle training on asthmatic patients was our aim in this study.

**Method:** This systematic review complied with PRISMA recommendation. As part of an extensive literature search, we searched the electronic databases PubMed, Scopus, and Web of Science for articles published between 2000 and 2017.

**Result and conclusion:** Six publications were considered in this analysis; one of them were quasi-experimental studies and five was a randomized controlled trial. Exercises and inspiratory muscle training are among the interventions employed. Exercise capacity, inspiratory muscle strength, exertional dyspnea, ER visits, asthma symptoms,  $\beta_2$ -agonist intake, IMS, and lung function were the outcome measures that were employed. An IMT program is a helpful intervention for improving IMS in people with asthma, according to this thorough evaluation. It also suggests that this intervention may have no adverse consequences and instead have a favorable effect on IMS, the requirement for rescue medication, and exertional dyspnea. Regarding how IMT affects exercise ability, hospital admissions, and health related QoL, the findings are contradicting. It had no effect on lung function or expiratory muscle strength.

**Keywords:** Respiratory muscle training; Exercise; Asthma; Treatment; Lung function

### 1. Introduction

One of the most common chronic respiratory conditions in the world is asthma (1). It is seen as a significant social and health issue and is linked to a significant financial burden, accounting for between 1% and 2% of all sanitary expenditures in developed nations (2). Chronic airway inflammation, which is typified by a history of respiratory symptoms like wheezing, shortness of breath, chest tightness, and coughing that fluctuate in duration and severity, together with variable expiratory airflow restriction, which may eventually become persistent, is what defines asthma. Some people with asthma may also have respiratory muscle dysfunction in addition to these symptoms (3).

This mechanical disadvantage of the diaphragm can cause dyspnea and greater strain for the inspiratory muscles, particularly during exercise when dynamic hyperinflation may develop (3). Furthermore, some research has


\* Corresponding author:

demonstrated that steroid-induced myopathy or muscular weakening might result from large dosages of systemic corticosteroids given to asthmatic patients (4). Additionally, people with mild stable asthma have been shown to have thoraco-abdominal asynchrony with moderate exercise. Because it is non-invasive and volitional, the evaluation of maximum respiratory pressures is frequently used in clinical practice to detect respiratory muscle weakness. When the readings fall between 65% and 80% of the anticipated range, the maximal inspiratory pressure (P<sub>Imax</sub>) and maximal expiratory pressures (P<sub>Emax</sub>) are deemed lowered (5).

When managing asthma over the long term, respiratory muscle dysfunction must be taken into account. Comprehensive programs incorporating education, breathing exercises, and exercise training have been emphasized as adjuvant therapy to asthma pharmaceutical treatment (6), in accordance with the chronic care paradigm and multidisciplinary approach. Nevertheless, these regimens have not consistently incorporated respiratory muscle training (RMT). RMT has demonstrated efficacy in treating patients with various respiratory deficits and chronic obstructive pulmonary disease (COPD) (7). RMT's usefulness in treating asthmatics is yet unknown, though. Our goal in this study was to examine how respiratory muscle training affects asthmatic patients.

## 2. Method

The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guideline was followed in the reporting of this systematic review (8).



**Figure 1** PRISMA consort chart

We looked through the electronic databases PubMed, Scopus, and Web of Science for papers published between 2000 and 2017 as part of a comprehensive literature search. Weekly automated updates that were obtained from these databases became the subject of additional investigation. A manual search of the major publications, earlier systematic reviews, and reference lists of the included studies was conducted in addition to the computerized check. search terms include; asthma, bronchoconstriction, bronchial spasm, breathing exercises, inspiratory muscle training, respiratory muscle training, expiratory muscle training, maximal voluntary ventilation, maximal respiratory pressures, dry powder inhalers, rescue medication, metered dose inhalers, respiratory function test, dyspnoea, adverse effects, exercise tolerance.

Quasi-experimental studies and RCTs involving a sample of asthma patients, irrespective of severity, were included in this systematic review. Any intervention that provides an external load to the airways using external device to increase IMS or endurance is referred to be an IMT or EMT program, and studies have to employ it as a stand-alone intervention.

Four writers individually examined the publications to find pertinent works based on the title and abstract after eliminating duplicate research. In the event of a dispute, the relevant author was consulted. Every article's citation, research design, participants, asthma severity, sample size, treatments, results, and conclusions were extracted using a preplanned process.

### 3. Result and discussion

In this study we included 6 articles, 5 were randomized controlled trials, and one study was n Quasi experimental. Interventions used include; inspiratory muscle training, and exercises. Outcome measures used were (Exercise capacity, inspiratory muscle strength (IMS), Exertional dyspnea, Emergency department visits, asthma symptoms, consumption of  $\beta$ 2-agonist, IMS and lung function (Table 1).

This comprehensive review shown that an IMT program is a useful intervention for enhancing IMS in individuals with asthma. It also raises the possibility that this intervention may positively impact IMS, the need for rescue medicine, and exertional dyspnea while having no negative side effects. The data are conflicting on the effects of IMT for exercise capacity, hospital admissions, and HRQoL. It did not increase expiratory muscular strength or lung function.

The present systematic review includes additional outcomes, such as exercise capacity, health related QoL, and respiratory muscle endurance, than the review published in 2013 (9). Higher PImax and PEmax were seen in the expiratory muscle training (EMT) group compared to the control group in a systematic evaluation of COPD patients, and these effects were more pronounced when IMT and EMT were combined. Moreover, there was a correlation found between increased exacerbations, hospital admissions, and death with weakening of the expiratory muscles (10). Thus, we think that more research is necessary to examine the potential advantages of EMT in conjunction with IMT for asthma.

When compared to no intervention or instructional programs, IMT can cause an increase in PImax in individuals with asthma. The majority of research conducted on chronic respiratory disorders used sham-IMT with training loads  $\leq$ 15% PImax, which did not result in changes to strength (6). Therefore, we think that the control group's inspiratory muscular strength was likely increased by a load of 20% PImax (11).

The primary conclusions of the TURNER et al. study are that asthmatic patients undergoing six weeks of IMT get a decreased oxygen consumption and the experience of dyspnea during exercise; enhanced IMS and demonstrated reduction of exercise-induced inspiratory muscle fatigue; and raised Tlim during constant-power exercise.

Consistent with the findings of TURNER et al. study, impaired IMS prior to exercise has been demonstrated to exacerbate exertional dyspnea in asthmatic individuals (12) and may also decrease exercise tolerance in these individuals (9). Additionally, IMT has been demonstrated to decrease the perception of exercise dyspnea in healthy individuals (13) and during pressurized threshold inspiratory loading in asthmatics (12). One important indicator of dyspnea during bronchoconstriction is the increased inspiratory muscle effort linked to increased airway resistance and dynamic hyperinflation (14) It might be caused by a variety of things, such as the recruitment of extra accessory respiratory muscles, a decrease in IMS because to their shorter operational duration, and an increase in inspiratory activity during expiration (15).

According to the TURNER et al. research, the decrease in dyspnea perception that they observed was probably caused by an IMS improvement following IMT rather than by modifications to lung function. Furthermore, literature indicates

a negative correlation between changes in IMS and the severity of dyspnea during an inspiratory loading test after IMT (16). To be more precise, after inspiratory muscle training, a lower proportion of maximum force-generating capacity would be needed to provide the necessary pressure for a given volume change. This would lower the central motor demand and, consequently, the perception of effort (17).

Aerobic training (18) and the injection of bronchodilators (19) have been demonstrated to diminish exertional dyspnea, one of the symptoms of asthma that is linked to poor exercise tolerance (20). TURNER et al. study conclude that, among recreationally active persons with moderate asthma, IMT can lessen the sense of dyspnea and increase exercise tolerance. This discovery raises the possibility of IMT serving as a supplemental intervention for this population. It should be highlighted, nonetheless, that for the tiny percentage of asthmatics who have a poor perception of dyspnea, a decrease in that perception might be harmful as it could lead to an under estimation of the severity of asthmatic exacerbations.

Asthma sufferers may have decreased exercise tolerance in correlation with expiratory flow restriction and dynamic hyperinflation during exercise (19). TURNER et al. study participants showed a progressive increase in end-inspiratory lung volume (EELV) as exercise duration increased; however, those in the IMT group exhibited a tendency toward a reduction in the degree of hyperinflation during the first five minutes of exercise following IMT, as well as a significant attenuation at that point. Following intervention, there was a comparable trend for EILV to decrease in the IMT group. According to earlier studies, a rise in EELV during intentional hyperpnea raised the oxygen cost of ventilation and the elastic work of breathing (21).

The findings of the Lima et al. study are consistent with those of studies involving adults with obstructive illnesses (16,22), demonstrating that following particular IMT, IMS improved, leading to a corresponding improvement in clinical status. An evaluation of a respiratory treatment program in eight distinct locations was conducted by a multicenter, randomized, cross-over controlled research (23). In addition to breathing exercises and instructional materials, the program featured IMT, relaxation methods, bronchial hygiene practices, and leisure activities. Exercise tolerance and patient quality of life were shown to have significantly improved.

**Table 1** Characteristics and main findings of the included studies

Citation	Study participants	Design	Outcomes measures	Intervention	Findings
Delgado et al. (24)	Controlled asthma	RCT	Exercise capacity and IMS	Inspiratory muscle training (IMT)	Intervention group PI <sub>max</sub> (cmH <sub>2</sub> O): pre vs post ( $p < 0.05$ ) 6MWT (m): pre vs post ( $p < 0.05$ ) No changes in pulmonary function:
Turner et al. (25)	Mild to moderate asthma	Quasi experimental	Exertional dyspnea and IMS	IMT	Following the intervention, there was no discernible difference in Time to the limit of exercise tolerance (Tlim), respiratory muscle exhaustion, or the placebo group's experience of dyspnea. A substantial 16% decrease in dyspnea during exercise was also observed. Despite the extended Tlim, the exercise-induced decline in intervention group was reduced from 10% before to IMT to 6% following IMT. The levels of pulmonary function in the placebo and IMT groups did not alter. IMT improves exercise tolerance, lessens the sensation of dyspnea, and attenuates inspiratory muscle fatigue. According to these results, IMT could be a useful addition to asthma treatment that helps

					increase this group's engagement and adherence to exercise regimens. Breathlessness, however, is also a significant indicator of bronchoconstriction, therefore if this symptom is unusually low, care should be taken.
Lima et al. (26)	Uncontrolled asthma	RCT	Emergency department visits, asthma symptoms and IMS	Breathing exercises, IMT	In addition to enhancing peak expiratory flow and severity factors, programs using IMT and respiratory workouts can boost the mechanical efficiency of the respiratory muscles.
Sampaio et al. (27)	Clinical diagnostic of asthma	RCT	IMS	IMT	In the intervention group PI <sub>max</sub> : Pre vs Post; $p < 0.05$ PE <sub>max</sub> : Pre vs Post; $p < 0.05$ No significant changes in PE <sub>max</sub> and PI <sub>max</sub>
Weiner et al. (16)	Mild to moderate asthma	RCT	Consumption of $\beta_2$ -agonist, IMS and lung function	IMT	The mean daily consumption of beta2-agonists and the baseline maximum inspiratory pressure did not exhibit a strong association with each other, nor with the perception of dyspnea (POD). Nonetheless, a noteworthy association was seen between the POD and the average daily intake of beta2-agonists. The decline in POD and the reduction in beta2-agonist intake were strongly linked with the increase in IMS following inspiratory muscle training.
Weiner et al. (28)	Mild to moderate asthma	RCT	$\beta_2$ -agonist consumption, IMS, and lung function.	IMT	In the female individuals, mean daily 2-agonist intake and POD were considerably greater, whereas baseline PI <sub>max</sub> was significantly lower. By the conclusion of the twentieth training week, PI <sub>max</sub> was on par with the male participants. A statistically significant reduction in mean daily 2-agonist usage and POD, comparable to that observed in male patients, was linked to an increase in PI <sub>max</sub> .

#### 4. Conclusion

This systematic review found that patients with asthma who participated in an IMT program saw a substantial improvement in inspiratory muscle strength but no change in expiratory muscle strength or lung function. Furthermore, our research reaffirms the applicability of the training's dose-response principle and suggests that this intervention may lessen the need for rescue medicine and the perception of dyspnea.

#### Compliance with ethical standards

*Disclosure of conflict of interest*

No conflict of interest to be disclosed.

## References

- [1] Soriano JB, Abajobir AA, Abate KH, Abera SF, Agrawal A, Ahmed MB, et al. Global, regional, and national deaths, prevalence, disability-adjusted life years, and years lived with disability for chronic obstructive pulmonary disease and asthma, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet Respir Med* [Internet]. 2017 Sep;5(9):691–706. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S221326001730293X>
- [2] Ehteshami-Afshar S, Fitzgerald JM, Doyle-Waters MM, Sadatsafavi M. The global economic burden of asthma and chronic obstructive pulmonary disease. *Int J Tuberc Lung Dis* [Internet]. 2016 Jan 1;20(1):11–23. Available from: <http://openurl.ingenta.com/content/xref?genre=article&issn=1027-3719&volume=20&issue=1&spage=11>
- [3] McConnell AK. The role of inspiratory muscle function and training in the genesis of dyspnoea in asthma and COPD. *Prim Care Respir J* [Internet]. 2005 Aug 1;14(4):186–94. Available from: <https://www.nature.com/articles/pcrj2005038>
- [4] Decramer M, Lacquet LM, Fagard R, Rogiers P. Corticosteroids contribute to muscle weakness in chronic airflow obstruction. *Am J Respir Crit Care Med* [Internet]. 1994 Jul;150(1):11–6. Available from: <https://www.atsjournals.org/doi/10.1164/ajrccm.150.1.8025735>
- [5] Barreiro E, Bustamante V, Cejudo P, Gálvez JB, Gea J, de Lucas P, et al. Normativa SEPAR sobre disfunción muscular de los pacientes con enfermedad pulmonar obstructiva crónica. *Arch Bronconeumol* [Internet]. 2015 Aug;51(8):384–95. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0300289615001568>
- [6] Shei RJ, Paris HLR, Wilhite DP, Chapman RF, Mickleborough TD. The role of inspiratory muscle training in the management of asthma and exercise-induced bronchoconstriction. *Phys Sportsmed* [Internet]. 2016 Oct 1;44(4):327–34. Available from: <https://www.tandfonline.com/doi/full/10.1080/00913847.2016.1176546>
- [7] Ramsook AH, Molgat-Seon Y, Schaeffer MR, Wilkie SS, Camp PG, Reid WD, et al. Effects of inspiratory muscle training on respiratory muscle electromyography and dyspnea during exercise in healthy men. *J Appl Physiol* [Internet]. 2017 May 1;122(5):1267–75. Available from: <https://www.physiology.org/doi/10.1152/japplphysiol.00046.2017>
- [8] Hutton B, Catalá-López F, Moher D. La extensión de la declaración PRISMA para revisiones sistemáticas que incorporan metaanálisis en red: PRISMA-NMA. *Med Clin (Barc)* [Internet]. 2016 Sep;147(6):262–6. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0025775316001512>
- [9] Silva IS, Fregonezi GA, Dias F AL, Ribeiro CT, Guerra RO, Ferreira GM. Inspiratory muscle training for asthma. *Cochrane Database Syst Rev* [Internet]. 2013 Sep 8; Available from: <https://doi.wiley.com/10.1002/14651858.CD003792.pub2>
- [10] Neves LF, Reis MH, Plentz RD, Matte DL, Coronel CC, Sbruzzi G. Expiratory and Expiratory Plus Inspiratory Muscle Training Improves Respiratory Muscle Strength in Subjects With COPD: Systematic Review. *Respir Care* [Internet]. 2014 Sep;59(9):1381–8. Available from: <http://rc.rcjournal.com/lookup/doi/10.4187/respcare.02793>
- [11] McConnell A, Caine M, Donovan K, Toogood A, Miller M. Inspiratory Muscle Training Improves Lung Function and Reduces Exertional Dyspnoea in Mild/Moderate asthmatics. *Clin Sci* [Internet]. 1998 Aug 1;95(s39):4P-4P. Available from: <https://portlandpress.com/clinsci/article/95/s39/4P/107429/Inspiratory-Muscle-Training-Improves-Lung-Function>
- [12] Killian KJ, Summers E, Watson RM, O'Byrne PM, Jones NL, Campbell EJ. Factors contributing to dyspnoea during bronchoconstriction and exercise in asthmatic subjects. *Eur Respir J* [Internet]. 1993 Jul;6(7):1004–10. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/8370426>
- [13] Romer LM, McConnell AK, Jones DA. Effects of inspiratory muscle training on time-trial performance in trained cyclists. *J Sports Sci* [Internet]. 2002 Jan 9;20(7):547–90. Available from: <https://www.tandfonline.com/doi/full/10.1080/02640410276000053>
- [14] Lougheed MD, Fisher T, O'Donnell DE. Dynamic Hyperinflation During Bronchoconstriction in Asthma. *Chest* [Internet]. 2006 Oct;130(4):1072–81. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0012369215511422>

[15] Martin J, Powell E, Shore S, Emrich J, Engel LA. The Role of Respiratory Muscles in the Hyperinflation of Bronchial Asthma 1– 3. *Am Rev Respir Dis* [Internet]. 1980 Mar;121(3):441–7. Available from: <http://www.atsjournals.org/doi/10.1164/arrd.1980.121.3.441>

[16] Weiner P, Magadle R, Beckerman M, Berar-Yanay N. The Relationship among the Inspiratory Muscle Strength, the Perception of Dyspnea and Inhaled Beta 2 -Agonists in Patients with Asthma. *Can Respir J* [Internet]. 2002 Jan 3;9(5):307–12. Available from: <https://onlinelibrary.wiley.com/doi/10.1155/2002/746808>

[17] el-Manshawi A, Killian KJ, Summers E, Jones NL. Breathlessness during exercise with and without resistive loading. *J Appl Physiol* [Internet]. 1986 Sep 1;61(3):896–905. Available from: <https://www.physiology.org/doi/10.1152/jappl.1986.61.3.896>

[18] Hallstrand TS, Bates PW, Schoene RB. Aerobic Conditioning in Mild Asthma Decreases the Hyperpnea of Exercise and Improves Exercise and Ventilatory Capacity. *Chest* [Internet]. 2000 Nov;118(5):1460–9. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0012369215512257>

[19] Kosmas EN. Exercise-induced flow limitation, dynamic hyperinflation and exercise capacity in patients with bronchial asthma. *Eur Respir J* [Internet]. 2004 Sep 1;24(3):378–84. Available from: <http://erj.ersjournals.com/cgi/doi/10.1183/09031936.04.00113003>

[20] O'DONNELL DE, REVILL SM, WEBB KA. Dynamic Hyperinflation and Exercise Intolerance in Chronic Obstructive Pulmonary Disease. *Am J Respir Crit Care Med* [Internet]. 2001 Sep 1;164(5):770–7. Available from: <https://www.atsjournals.org/doi/10.1164/ajrccm.164.5.2012122>

[21] Collett PW, Perry C, Engel LA. Pressure-time product, flow, and oxygen cost of resistive breathing in humans. *J Appl Physiol* [Internet]. 1985 Apr 1;58(4):1263–72. Available from: <https://www.physiology.org/doi/10.1152/jappl.1985.58.4.1263>

[22] Weiner P, Berar-Yanay N, Davidovich A, Magadle R, Weiner M. Specific Inspiratory Muscle Training in Patients With Mild Asthma With High Consumption of Inhaled  $\beta$ 2-Agonists. *Chest* [Internet]. 2000 Mar;117(3):722–7. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S001236921532729X>

[23] Cambach W, Chadwick-Straver R, Wagenaar R, van Keimpema A, Kemper H. The effects of a community-based pulmonary rehabilitation programme on exercise tolerance and quality of life: a randomized controlled trial. *Eur Respir J* [Internet]. 1997 Jan 1;10(1):104–13. Available from: <http://erj.ersjournals.com/lookup/doi/10.1183/09031936.97.10010104>

[24] Delgado, R., Silva, I., Silva, T. E., Fernandes, T., Lima Filho, N., Oliveira, V. H., ... & Ferreira, G. (2014). Inspiratory muscle training for asthma—A randomized controlled pilot study. *European Respiratory Journal*, 44(Suppl 58)..

[25] TURNER LA, MICKLEBOROUGH TD, MCCONNELL AK, STAGER JM, TECKLENBURG-LUND S, LINDLEY MR. Effect of Inspiratory Muscle Training on Exercise Tolerance in Asthmatic Individuals. *Med Sci Sport Exerc* [Internet]. 2011 Nov;43(11):2031–8. Available from: <https://journals.lww.com/00005768-201111000-00003>

[26] Lima EVNCL, Lima WL, Nobre A, Santos AM dos, Brito LMO, Costa M do R da SR. Treinamento muscular inspiratório e exercícios respiratórios em crianças asmáticas. *J Bras Pneumol* [Internet]. 2008 Aug;34(8):552–8. Available from: [http://www.scielo.br/scielo.php?script=sci\\_arttext&pid=S1806-37132008000800003&lng=pt&tlang=pt](http://www.scielo.br/scielo.php?script=sci_arttext&pid=S1806-37132008000800003&lng=pt&tlang=pt)

[27] Sampaio, L. M. M., Jamami, M., Pires, V. A., Silva, A. B., & Costa, D. (2002). Respiratory muscle strength in asthmatic patient submitted by respiratory muscle training and physical training. *Physiother Mag Univ Sao Paolo*, 9(2), 43–48..

[28] Weiner P, Magadle R, Massarwa F, Beckerman M, Berar-Yanay N. Influence of Gender and Inspiratory Muscle Training on the Perception of Dyspnea in Patients With Asthma. *Chest* [Internet]. 2002 Jul;122(1):197–201. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0012369216462999>